COVID-19 Vaccines & Outpatient Therapeutics

A quick reference guide for healthcare providers

COVID-19 Vaccines

Pfizer-BioNTech (Comirnaty): for ages 5 years and older	**Moderna (mNexspike): for ages 12 years and older**
Moderna (Spikevax): for ages 6 months and older	**Novavax (Nuvaxovid): for ages 12 years and older**

COVID-19 Preventive Therapeutics

Post-exposure prophylaxis

- **Ensitrelvir:** a preventive oral antiviral for individuals following contact with someone who has COVID-19. This is indicated for people ages 12 years and older. This should be taken within 72 hours of contact.

Pre-exposure prophylaxis for individuals moderately to severely immunocompromised

- **Pemivibart:** a preventive monoclonal antibody administered as an IV infusion. This is indicated for people ages 12 years or older and weighing at least 88 pounds and should be used in conjunction with vaccination.

COVID-19 Antiviral Treatments

- **Nirmatrelvir with Ritonavir:** an oral medication for ages 12 and older to be taken within 5 days of symptom onset.
- **Remdesivir:** an antiviral administered through IV infusion for three consecutive days for all ages. This treatment should begin within 7 days of symptom onset.
- **Molnupiravir:** an oral medication for ages 18 and older to be taken within 5 days of symptom onset.

Keep the latest healthcare provider resources at your fingertips with the American Lung Association's COVID-19 Resource Hub. Visit **Lung.org/hcp-covid**

Support for this education provided by Shionogi | September 2026